arXiv:1610.04127v1 [math.AP] 13 Oct 2016

THE MAZ'YA-SHAPOSHNIKOVA LIMIT IN THE MAGNETIC SETTING

ANDREA PINAMONTI, MARCO SQUASSINA, AND EUGENIO VECCHI

Abstract. We prove a magnetic version of the Maz'ya-Shaposhnikova singular limit of nonlocal norms with vanishing fractional parameter. This complements a general convergence result recently obtained by authors when the parameter approaches one.

1. Introduction

About fifteen years ago, V. Maz'ya and T. Shaposhnikova proved that for any $n \geq 1$ and $p \in [1, \infty)$,

$$\lim_{s \searrow 0} s \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x) - u(y)|^p}{|x-y|^{n+ps}} dxdy = \frac{4\pi^{n/2}}{p\Gamma(n/2)} \|u\|^p_{L^p(\mathbb{R}^n)},$$

whenever $u \in D^{s,p}_0(\mathbb{R}^n)$ for some $s \in (0,1)$. Here Γ denotes the Gamma function and the space $D^{s,p}_0(\mathbb{R}^n)$ is the completion of $C^\infty_c(\mathbb{R}^n)$ with respect to the Gagliardo norm

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x) - u(y)|^p}{|x-y|^{n+ps}} dxdy.$$

Their motivation was basically that of complementing a previous result by Bourgain-Brezis-Mironescu [4,5] providing new characterizations for functions in the Sobolev space $W^{1,p}(\Omega)$. Precisely, if $\Omega \subset \mathbb{R}^n$ is a smooth bounded domain, then for any $W^{1,p}(\Omega)$ there holds

$$\lim_{s \nearrow 1} (1-s) \int_\Omega \int_\Omega \frac{|u(x) - u(y)|^p}{|x-y|^{n+ps}} dxdy = Q_{p,n} \int_\Omega |\nabla u|^p dx,$$

where $Q_{p,n}$ is defined by

$$Q_{p,n} = \frac{1}{p} \int_{\mathbb{S}^{n-1}} |\boldsymbol{\omega} \cdot h|^p d\mathcal{H}^{n-1}(h), \tag{1.1}$$

being $\mathbb{S}^{n-1}$ the unit sphere in $\mathbb{R}^n$ and $\boldsymbol{\omega}$ an arbitrary unit vector of $\mathbb{R}^n$. The above singular limits are natural and also admit a physical relevance in the framework of the theory of Levy processes. Also, there is a developed theory of fractional s-perimeters [6] and there have been several contributions concerning their asymptotic analysis in the limits $s \nearrow 1$ and $s \searrow 0$ [1,7,9].

One of the latest generalizations of this kind of convergence results appeared recently in [16] in the context of *magnetic Sobolev spaces* $W^{1,2}_A(\Omega)$, see [12]. In fact, a relevant role in the study of particles which interact with a magnetic field $B = \nabla \times A$, $A : \mathbb{R}^n \to \mathbb{R}^n$, is assumed by the *magnetic Laplacian* $(\nabla - \mathrm{i}A)^2$ [3,12,15], yielding to nonlinear Schrödinger equations of the type $-(\nabla - \mathrm{i}A)^2 u + u = f(u)$, which have been extensively studied (see [2] and the references therein). The operator is defined weakly as the differential of the energy

$$W^{1,2}_A(\Omega) \ni u \mapsto \int_\Omega |\nabla u - \mathrm{i}A(x)u|^2 dx.$$

If $A : \mathbb{R}^n \to \mathbb{R}^n$ is a smooth field and $s \in (0,1)$, a nonlocal magnetic counterpart of the magnetic laplacian,

$$(-\Delta)^s_A u(x) = c(n,s) \lim_{\varepsilon \searrow 0} \int_{B^c_\varepsilon(x)} \frac{u(x) - e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)} u(y)}{|x-y|^{n+2s}} dy,$$

2010 *Mathematics Subject Classification.* 49A50, 26A33, 82D99.

Key words and phrases. Fractional magnetic spaces, Maz'ya-Shaposhnikova limit, characterization of Sobolev spaces.

The authors are members of *Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni* (GNAMPA) of the *Istituto Nazionale di Alta Matematica* (INdAM). E.V. receives funding from the People Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme FP7/2007-2013/ under REA grant agreement No. 607643 (ERC Grant MaNET 'Metric Analysis for Emergent Technologies').

where $c(n,s)$ is a normalization constant which behaves as follows

$$\lim_{s\searrow 0}\frac{c(n,s)}{s}=\frac{\Gamma(n/2)}{\pi^{n/2}},\qquad \lim_{s\nearrow 1}\frac{c(n,s)}{1-s}=\frac{2n\Gamma(n/2)}{\pi^{n/2}}, \tag{1.2}$$

was introduced in [8, 11] for complex-valued functions, with motivations falling into the framework of the general theory of Lévy processes. Recently, the authors in [14] (see [16] for $p=2$) proved that if $A:\mathbb{R}^n\to\mathbb{R}^n$ is a C^2 vector field, then, for any $n\geq 1$, $p\in[1,\infty)$ and any Lipschitz bounded domain $\Omega\subset\mathbb{R}^n$

$$\lim_{s\nearrow 1}(1-s)\int_\Omega\int_\Omega\frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|_p^p}{|x-y|^{n+ps}}dxdy=Q_{p,n}\int_\Omega|\nabla u-\mathrm{i}A(x)u|_p^p\,dx, \tag{1.3}$$

for all $u\in W^{1,p}_A(\Omega)$, where $Q_{p,n}$ is as in (1.1) and $|z|_p:=\left(|(\Re z_1,\dots,\Re z_n)|^p+|(\Im z_1,\dots,\Im z_n)|^p\right)^{1/p}$. This has provided a new nonlocal characterization of the magnetic Sobolev spaces $W^{1,p}_A(\Omega)$.

The main goal of this paper is to complete the picture of [14] by providing a *magnetic counterpart* of the convergence result by *Maz'ya-Shaposhnikova* for vanishing fractional orders s, namely for $s\searrow 0$.

We consider a locally bounded vector potential field $A:\mathbb{R}^n\to\mathbb{R}^n$ and the space of complex valued functions $D^{s,p}_{A,0}(\mathbb{R}^n,\mathbb{C})$ defined as the completion of $C^\infty_c(\mathbb{R}^n,\mathbb{C})$ with respect to the norm

$$\|u\|_{D^{s,p}_{A,0}}=\left(\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+ps}}dxdy\right)^{1/p}.$$

By combining Lemma 2.1 and Lemma 2.3, we shall prove the following result.

Theorem 1.1 (Magnetic Maz'ya-Shaposhnikova)**.** *Let $n\geq 1$ and $p\in[1,\infty)$. Then for every*

$$u\in\bigcup_{0<s<1}D^{s,p}_{A,0}(\mathbb{R}^n,\mathbb{C}),$$

there holds

$$\lim_{s\searrow 0}s\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+ps}}dxdy=\frac{4\pi^{n/2}}{p\Gamma(n/2)}\|u\|^p_{L^p(\mathbb{R}^n)}.$$

In particular, while the singular limit as $s\nearrow 1$ generates the magnetic gradient $\nabla-\mathrm{i}A$, the limit for vanishing s tends to destroy the magnetic effects yielding the $L^p(\mathbb{R}^n)$-norm of the function u. We point out that, while in (1.3) the norm of complex numbers is $|\cdot|_p$, in Theorem 1.1 we use the usual norm $|\cdot|=|\cdot|_2$. In any case when $A=0$ and u is real-valued the formulas are all consistent with the classical statements. In the case $p=2$, combining the asymptotic formulas in (1.2) with Theorem 1.1 implies that

$$\frac{c(n,s)}{2}\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^2}{|x-y|^{n+2s}}dxdy\approx\|u\|^2_{L^2(\mathbb{R}^n)},\qquad\text{as } s\searrow 0,$$

for any $u\in D^{s,2}_0(\mathbb{R}^n)$ for some $s\in(0,1)$. Although the magnetic setting is mainly meaningful in the framework of nonlocal Schrödinger equations, we remark that for $E\subset\mathbb{R}^n$, if $E^c:=\mathbb{R}^n\setminus E$, the quantity

$$P_s(E;A):=\frac{1}{2}\int_E\int_E\frac{|1-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}|}{|x-y|^{n+s}}dxdy+\int_E\int_{E^c}\frac{1}{|x-y|^{n+s}}dxdy$$

plays the role of a nonlocal s-perimeter of E depending on A, which reduces for $A=0$ to the usual notion of fractional s-perimeter of $E\subset\mathbb{R}^n$. Then, if $\mathscr{L}^n(E)$ denotes the n-dimensional Lebesgue measure of $E\subset\mathbb{R}^n$, Theorem 1.1, applied with $p=1$ and $u(x)=\mathbf{1}_E(x)$, reads as

$$\lim_{s\searrow 0}sP_s(E,A)=\frac{4\pi^{n/2}}{\Gamma(n/2)}\mathscr{L}^n(E),$$

provided that $P_{s_0}(E,A)<+\infty$, for some $s_0\in(0,1)$.

2. Proof of the main result

The proof of Theorem 1.1 follows by combining Lemma 2.1 and Lemma 2.3 below.

Lemma 2.1 (Liminf inequality). *Let $n \geq 1$, $p \in [1,\infty)$ and let*

$$u \in \bigcup_{0<s<1} D^{s,p}_{A,0}(\mathbb{R}^n,\mathbb{C}).$$

Then

$$\liminf_{s\searrow 0} s \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+ps}}dxdy \geq \frac{4\pi^{n/2}}{p\Gamma(n/2)}\|u\|^p_{L^p(\mathbb{R}^n)}.$$

Proof. If

$$\liminf_{s\searrow 0} s \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+ps}}dxdy = \infty,$$

the assertion follows. Otherwise, there exists a sequence $\{s_k\}_{k\in\mathbb{N}} \subset (0,1)$ with $s_k \searrow 0$ and

$$\liminf_{s\searrow 0} s \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+ps}}dxdy = \lim_{k\to\infty} s_k \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+ps_k}}dxdy,$$

the limit being finite. For a.e. $x,y\in\mathbb{R}^n$ we have the Diamagnetic inequality (cf. [8, Remark 3.2])

$$||u(x)|-|u(y)|| \leq |u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|. \tag{2.1}$$

In particular, since $u\in D^{s_k,p}_{A,0}(\mathbb{R}^n,\mathbb{C})$, we have $|u|\in D^{s_k,p}_0(\mathbb{R}^n)$ and, for any $k\geq 1$,

$$s_k \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{||u(x)|-|u(y)||^p}{|x-y|^{n+ps_k}}dxdy \leq s_k \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+ps_k}}dxdy.$$

Taking the limit as $k\to\infty$ on both sides and invoking [13, Theorem 3] applied to $|u|$, yields

$$\frac{4\pi^{n/2}}{p\Gamma(\frac{n}{2})}\||u|\|^p_{L^p(\mathbb{R}^n)} \leq \lim_{k\to\infty} s_k \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+ps_k}}dxdy,$$

which concludes the proof. □

Remark 2.2 (Magnetic Hardy inequality). By combining the pointwise Diamagnetic inequality (2.1) with the fractional Hardy inequality [10], for $n > ps$ the following *magnetic Hardy inequality* holds: there exists a positive constant $\mathcal{H}_{n,s,p}$ such that

$$\int_{\mathbb{R}^n} \frac{|u(x)|^p}{|x|^{sp}}dx \leq \mathcal{H}_{n,s,p}\int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+ps}}dxdy, \tag{2.2}$$

for every $u\in D^{s,p}_{A,0}(\mathbb{R}^n,\mathbb{C})$. Similarly the following *magnetic Sobolev inequality* holds: there exists a positive constant $\mathcal{S}_{n,s,p}$ such that

$$\Big(\int_{\mathbb{R}^n} |u(x)|^{\frac{np}{n-sp}}dx\Big)^{\frac{n-sp}{n}} \leq \mathcal{S}_{n,s,p}\int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+ps}}dxdy,$$

for every $u\in D^{s,p}_{A,0}(\mathbb{R}^n,\mathbb{C})$.

Next we state a second lemma completing the proof of Theorem 1.1 when combined with Lemma 2.1.

Lemma 2.3 (Limsup inequality). *Let $n\geq 1$, $p\in[1,\infty)$ and let*

$$u \in \bigcup_{0<s<1} D^{s,p}_{A,0}(\mathbb{R}^n,\mathbb{C}).$$

Then

$$\limsup_{s\searrow 0} s \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+ps}}dxdy \leq \frac{4\pi^{n/2}}{p\Gamma(n/2)}\|u\|^p_{L^p(\mathbb{R}^n)}.$$

Proof. If $u \notin L^p(\mathbb{R}^n)$, there is nothing to prove. Hence, we may assume that $u \in L^p(\mathbb{R}^n)$. We observe that

$$\begin{aligned}
&s\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+ps}}dxdy\\
&= s\int_{\mathbb{R}^n}\int_{\{|x|\le|y|\le 2|x|\}}\frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+sp}}dxdy\\
&+ s\int_{\mathbb{R}^n}\int_{\{|y|\ge 2|x|\}}\frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+sp}}dxdy\\
&+ s\int_{\mathbb{R}^n}\int_{\{|x|\ge|y|\}}\frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+sp}}dxdy\\
&= 2s\int_{\mathbb{R}^n}\int_{\{|x|\le|y|\le 2|x|\}}\frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+sp}}dxdy\\
&+ 2s\int_{\mathbb{R}^n}\int_{\{|y|\ge 2|x|\}}\frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+sp}}dxdy,
\end{aligned}$$

where the last equality follows noticing that since $|e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}| = 1$ then

$$\begin{aligned}
\int_{\mathbb{R}^n}\int_{\{|x|\ge|y|\}}\frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+sp}}dxdy &= \int_{\mathbb{R}^n}\int_{\{|y|\ge|x|\}}\frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+sp}}dxdy\\
&= \int_{\mathbb{R}^n}\int_{\{|y|\ge 2|x|\}}\frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+sp}}dxdy\\
&+ \int_{\mathbb{R}^n}\int_{\{|x|\le|y|\le 2|x|\}}\frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+sp}}dxdy.
\end{aligned}$$

Using the triangle inequality for the L^p-norm on $\mathbb{R}^{2n}$ and recalling that $|e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}| = 1$, yields

$$\begin{aligned}
&s\int_{\mathbb{R}^n}\int_{\{|y|\ge 2|x|\}}\frac{|u(x)-e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)}u(y)|^p}{|x-y|^{n+sp}}dxdy\\
&\le\left\{\left(s\int_{\mathbb{R}^n}\int_{\{|y|\ge 2|x|\}}\frac{|u(x)|^p}{|x-y|^{n+sp}}dxdy\right)^{1/p}+\left(s\int_{\mathbb{R}^n}\int_{\{|y|\ge 2|x|\}}\frac{|u(y)|^p}{|x-y|^{n+sp}}dxdy\right)^{1/p}\right\}^p.
\end{aligned}$$

We claim that

$$\lim_{s\searrow 0} s\int_{\mathbb{R}^n}\int_{\{|y|\ge 2|x|\}}\frac{|u(y)|^p}{|x-y|^{n+sp}}dxdy = 0.$$

Observe that $2|x-y| \ge |y| + (|y|-2|x|)$. Then, if $|y| \ge 2|x|$ we get $2|x-y| \ge |y|$. Now, if $\mathscr{H}^{n-1}$ denotes the $(n-1)$-dimensional Haudorff measure, it follows that

$$\begin{aligned}
s^{1/p}\left(\int_{\mathbb{R}^n}\int_{\{|y|\ge 2|x|\}}\frac{|u(y)|^p}{|x-y|^{n+sp}}dxdy\right)^{1/p} &\le s^{1/p}\left(2^{n+sp}\int_{\mathbb{R}^n}\frac{|u(y)|^p}{|y|^{n+sp}}\Big(\int_{\{|x|\le|y|/2\}}dx\Big)dy\right)^{1/p}\\
&= 2^s\left(\frac{s}{n}\mathscr{H}^{n-1}(\mathbb{S}^{n-1})\right)^{1/p}\left(\int_{\mathbb{R}^n}\frac{|u(y)|^p}{|y|^{sp}}dy\right)^{1/p},
\end{aligned}$$

and the last term goes to zero as $s \searrow 0$. Notice that $y \mapsto |y|^{-s}u(y)$ remains bounded in $L^p(\mathbb{R}^n)$ as $s \searrow 0$ by the argument indicated here below. Observe now that, if $|y| \geq 2|x|$ we then get $|x-y| \geq |x|$ yielding

$$\left(s \int_{\mathbb{R}^n} \int_{\{|y|\geq 2|x|\}} \frac{|u(x)|^p}{|x-y|^{n+sp}} dxdy \right)^{1/p} \leq \left(s \int_{\mathbb{R}^n} \int_{\{|x-y|\geq |x|\}} \frac{|u(x)|^p}{|x-y|^{n+sp}} dxdy \right)^{1/p}$$
$$= \left(s \int_{\mathbb{R}^n} |u(x)|^p \int_{B(0,|x|)^c} \frac{dz}{|z|^{n+sp}} dx \right)^{1/p} = \frac{\mathscr{H}^{n-1}(\mathbb{S}^{n-1})^{1/p}}{p^{1/p}} \left(\int_{\mathbb{R}^n} \frac{|u(x)|^p}{|x|^{sp}} dx \right)^{1/p}.$$

Moreover $|x|^{-sp}|u(x)|^p = f_s(x) + g_s(x)$, where

$$f_s(x) := \frac{|u(x)|^p}{|x|^{sp}} \mathbf{1}_{B(0,1)}(x), \qquad g_s(x) := \frac{|u(x)|^p}{|x|^{sp}} \mathbf{1}_{B(0,1)^c}(x) \leq |u(x)|^p \mathbf{1}_{B(0,1)^c}(x) \in L^1(\mathbb{R}^n),$$

and $s \mapsto f_s$ is decreasing and, moreover, by the Hardy inequality (2.2) and the assumption on u, it follows that $f_{\tilde{s}} \in L^1(\mathbb{R}^n)$ for some $\tilde{s} \in (0,1)$. Hence, by monotone and dominated convergence, we conclude that

$$\limsup_{s \searrow 0} s \int_{\mathbb{R}^n} \int_{\{|y|\geq 2|x|\}} \frac{|u(x)|^p}{|x-y|^{n+sp}} dxdy \leq \frac{\mathscr{H}^{n-1}(\mathbb{S}^{n-1})}{p} \|u\|^p_{L^p(\mathbb{R}^n)} = \frac{2\pi^{n/2}}{p\Gamma(\frac{n}{2})} \|u\|^p_{L^p(\mathbb{R}^n)}.$$

Then, we conclude from the above inequalities that

$$\limsup_{s \searrow 0} 2s \int_{\mathbb{R}^n} \int_{\{|y|\geq 2|x|\}} \frac{|u(x) - e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)} u(y)|^p}{|x-y|^{n+sp}} dxdy \leq \frac{4\pi^{n/2}}{p\Gamma(\frac{n}{2})} \|u\|^p_{L^p(\mathbb{R}^n)}. \tag{2.3}$$

We claim that

$$\limsup_{s \searrow 0} 2s \int_{\mathbb{R}^n} \int_{\{|x|\leq |y|\leq 2|x|\}} \frac{|u(x) - e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)} u(y)|^p}{|x-y|^{n+sp}} \, dxdy = 0. \tag{2.4}$$

By assumption let $\tau \in (0,1)$ such that $u \in D^{\tau,p}_{A,0}(\mathbb{R}^n)$. Now let $N \geq 1$ and $s < \tau$. Then

$$2s \int_{\mathbb{R}^n} \int_{\{|x|\leq |y|\leq 2|x|\}} \frac{|u(x) - e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)} u(y)|^p}{|x-y|^{n+sp}} \, dxdy$$
$$= 2s \int_{\mathbb{R}^n} \int_{\substack{\{|x-y|\leq N\} \\ \{|x|\leq |y|\leq 2|x|\}}} \frac{|u(x) - e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)} u(y)|^p}{|x-y|^{n+sp}} \, dxdy$$
$$+ 2s \int_{\mathbb{R}^n} \int_{\substack{\{|x-y|> N\} \\ \{|x|\leq |y|\leq 2|x|\}}} \frac{|u(x) - e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)} u(y)|^p}{|x-y|^{n+sp}} \, dxdy =: \mathcal{I} + \mathcal{II}.$$

Let us consider $\mathcal{I}$ first. Since $|x-y| \leq N$, it holds that

$$\frac{1}{|x-y|^{n+sp}} = \frac{|x-y|^{p(\tau-s)}}{|x-y|^{n+\tau p}} \leq \frac{N^{p(\tau-s)}}{|x-y|^{n+\tau p}}.$$

Therefore $\mathcal{I}$ goes to zero as $s \searrow$, since

$$\mathcal{I} \leq 2sN^{p(\tau-s)} \int_{\mathbb{R}^n} \int_{\substack{\{|x-y|\leq N\} \\ \{|x|\leq |y|\leq 2|x|\}}} \frac{|u(x) - e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)} u(y)|^p}{|x-y|^{n+\tau p}} \, dxdy.$$

Let us now move to $\mathcal{II}$. Since $|u(x) - e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)} u(y)|^p \leq 2^{p-1} \left(|u(x)|^p + |u(y)|^p\right)$, we get

$$\mathcal{II} \leq 2^p s \int_{\mathbb{R}^n} \int_{\substack{\{|x-y|\geq N\} \\ \{|x|\leq |y|\leq 2|x|\}}} \frac{|u(x)|^p}{|x-y|^{n+sp}} dxdy + 2^p s \int_{\mathbb{R}^n} \int_{\substack{\{|x-y|\geq N\} \\ \{|x|\leq |y|\leq 2|x|\}}} \frac{|u(y)|^p}{|x-y|^{n+sp}} dxdy =: \mathcal{II}' + \mathcal{II}''.$$

Regarding $\mathcal{II}'$, since $|x-y| \geq N$ and $|y| \leq 2|x|$, it holds

$$N \leq |x-y| \leq |x| + |y| \leq 3|x|,$$

which implies that $|x| \geq \frac{N}{3}$. In particular, this also implies that

$$\mathcal{II}' \leq 2^p s \int_{\{|x|\geq N/3\}} \left(\int_{\{|x-y|\geq N\}} \frac{|u(x)|^p}{|x-y|^{n+sp}} dy \right) dx \leq C(n,p) \int_{\{|x|\geq N/3\}} |u(x)|^p \, dx.$$

For $\mathcal{II}''$, since as before $|x-y| \geq N$ and $|x| \leq |y|$, we have

$$N \leq |x-y| \leq |x| + |y| \leq 2|y|,$$

which implies $|y| \geq \dfrac{N}{2} \geq \dfrac{N}{3}$. Therefore, we get

$$\mathcal{II}'' \leq 2^p s \int_{\{|y|\geq N/3\}} |u(y)|^p \left(\int_{\{|z|\geq N\}} \frac{1}{|z|^{n+sp}} dz \right) dy \leq C(n,p) \int_{\{|y|\geq N/3\}} |u(y)|^p \, dy.$$

Combining the estimates for $\mathcal{II}'$ and $\mathcal{II}''$, we get

$$\mathcal{II} \leq C(n,p) \int_{\{|x|\geq N/3\}} |u(x)|^p dx,$$

which is a bound independent of s. Now, going back to

$$\limsup_{s \searrow 0} 2s \int_{\mathbb{R}^n} \int_{\{|x|<|y|<2|x|\}} \frac{|u(x) - e^{\mathrm{i}(x-y)\cdot A\left(\frac{x+y}{2}\right)} u(y)|^p}{|x-y|^{n+sp}} \, dxdy \leq 2C(n,p) \|u\|^p_{L^p(B(0,N/3)^c)},$$

and (2.4) follows letting $N \to \infty$, since $u \in L^p(\mathbb{R}^n)$. Collecting (2.3) and (2.4), the assertion follows. □

References

[1] L. Ambrosio, G. De Philippis, L. Martinazzi, *Γ−convergence of nonlocal perimeter functionals,* Manuscripta Math. **134** (2011), 377–403. 1

[2] G. Arioli, A. Szulkin, *A semilinear Schrödinger equation in the presence of a magnetic field*, Arch. Ration. Mech. Anal. **170** (2003), 277–295. 1

[3] J. Avron, I. Herbst, B. Simon, *Schrödinger operators with magnetic fields. I. General interactions*, Duke Math. J. **45** (1978), 847–883. 1

[4] J. Bourgain, H. Brezis, P. Mironescu, *Another look at Sobolev spaces*, in *Optimal Control and Partial Differential Equations. A Volume in Honor of Professor Alain Bensoussan's 60th Birthday* (eds. J. L. Menaldi, E. Rofman and A. Sulem), IOS Press, Amsterdam, 2001, 439–455. 1

[5] J. Bourgain, H. Brezis, P. Mironescu, *Limiting embedding theorems for $W^{s,p}$ when $s \uparrow 1$ and applications*, J. Anal. Math. **87** (2002), 77–101. 1

[6] L. Caffarelli, J. -M. Roquejoffre, O. Savin, *Nonlocal minimal surfaces*, Comm. Pure Appl. Math. **63** (2010), 1111–1144. 1

[7] L. Caffarelli, E. Valdinoci, *Regularity properties of nonlocal minimal surfaces via limiting arguments*, Adv. Math. **248** (2013), 843–871. 1

[8] P. d'Avenia, M. Squassina, *Ground states for fractional magnetic operators*, preprint, http://arxiv.org/abs/1601.04230 2, 3

[9] S. Dipierro, A. Figalli, G. Palatucci, E. Valdinoci, *Asymptotics of the s-perimeter as $s \to 0$,* Discrete Contin. Dyn. Syst. **33** (2013), 2777–2790. 1

[10] R.L. Frank, R. Seiringer, *Non-linear ground state representation and sharp Hardy inequalities*, J. Funct. Anal. **255** (2008), 3407–3430. 3

[11] T. Ichinose, *Magnetic relativistic Schrödinger operators and imaginary-time path integrals*, Mathematical physics, spectral theory and stochastic analysis, 247–297, Oper. Theory Adv. Appl. **232**, Birkhäuser/Springer, Basel, 2013. 2

[12] E. Lieb and M. Loss, Analysis, Graduate Studies in Mathematics **14**, 2001. 1

[13] V. Maz'ya and T. Shaposhnikova, *On the Bourgain, Brezis, and Mironescu theorem concerning limiting embeddings of fractional Sobolev spaces*, J. Funct. Anal. **195** (2002), 230–238. 3

[14] A. Pinamonti, M. Squassina, E. Vecchi, *Magnetic BV functions and the Bourgain-Brezis-Mironescu formula*, preprint, https://arxiv.org/abs/1609.09714 2

[15] M. Reed, B. Simon, Methods of modern mathematical physics, I, Functional analysis, Academic Press, Inc., New York, 1980 1

[16] M. Squassina, B. Volzone, *Bourgain-Brezis-Mironescu formula for magnetic operators,* C. R. Math. Acad. Sci. Paris **354** (2016), 825–831. 1, 2

(A. Pinamonti) Dipartimento di Matematica
Università degli Studi di Padova, Via Trieste 63, 35121 Padova, Italy
E-mail address: Pinamont@math.unipd.it

(M. Squassina) Dipartimento di Matematica e Fisica
Università Cattolica del Sacro Cuore, Via dei Musei 41, I-25121 Brescia, Italy
E-mail address: marco.squassina@dmf.unicatt.it

(E. Vecchi) Dipartimento di Matematica
Università di Bologna, Piazza di Porta S. Donato 5, 40126, Bologna, Italy
E-mail address: eugenio.vecchi2@unibo.it